

wave-form analysis to compare with our laboratory data. Although these data may not be representative of all the reported seismic observations, they are probably the most sensitive to the features that we utilize and serve to illustrate the utility of our data in interpreting the seismic structure of the mantle.

Let us compare the differences in acoustic velocities between the olivine and modified spinel phase as deduced in this report with reported velocity discontinuities at 400 km in the mantle. The seismic compressional discontinuity of 0.43 km/sec (10) is about 40 percent of the difference between the two phases (1.06 km/sec). The seismic shear discontinuity of 0.23 km/sec (11) is about 35 percent of this difference (0.71 km/sec). This comparison suggests an olivine content of the mantle of 40 percent by volume, with the other 60 percent being composed of minerals that do not undergo phase transformations at this depth. This amount of olivine is low compared to that of most pyrolite mantle models, which generally require that olivine represent over 60 percent of the total volume (4). The pressure dependence of the velocity difference between the olivine and the beta phase is probably the greatest experimental uncertainty, and the magnitude of the velocity discontinuity is the greatest observational uncertainty. If the pressure derivatives of the bulk and shear moduli for the beta phase are approximately 3.9 and 1.1, then the observed discontinuity is consistent with a pyrolite composition. This compares to values of 5.3 and 1.8 for olivine. Although these inferred values are lower than expected, they are not sufficiently unreasonable to warrant rejecting the pyrolite model. We conclude that, although pyrolite is still an acceptable mantle model, a model with less olivine is preferable.

HIROSHI SAWAMOTO*
DONALD J. WEIDNER
SATOSHI SASAKI†
MINEO KUMAZAWA‡

Department of Earth and Space
Sciences, State University of
New York, Stony Brook 11794

References and Notes

1. K. Bullen, *An Introduction to the Theory of Seismology* (Cambridge Univ. Press, Cambridge, ed. 3, 1963).
2. F. Birch, *J. Geophys. Res.* **57**, 227 (1952).
3. L. R. Johnson, *ibid.* **72**, 6309 (1967).
4. A. E. Ringwood, *Composition and Petrology of the Earth's Mantle* (McGraw-Hill, New York, 1975).
5. D. J. Weidner and H. R. Carleton, *J. Geophys. Res.* **82**, 1334 (1977).
6. D. J. Weidner, H. Wang, J. Ito, *Phys. Earth Planet. Inter.* **17**, 7 (1978); D. J. Weidner and M. T. Vaughan, in *High-Pressure Science and Technology*, D. Timmerhaus and M. S. Barber, Eds. (Plenum, New York, 1979), vol. 2, p. 85.
7. H. Horiuchi and H. Sawamoto, *Am. Mineral.* **66**, 568 (1981).
8. H. Sawamoto, H. Horiuchi, M. Tokonami, M. Kumazawa, paper presented at the U.S.-Japan Seminar on High-Pressure Research: Applications in Geophysics, Hakone, Japan, 1981 (program with abstracts, p. 68).
9. M. Kumazawa and O. L. Anderson, *J. Geophys. Res.* **74**, 5961 (1969).
10. L. J. Burdick, *ibid.* **86**, 5926 (1981).
11. S. P. Grand and D. V. Helmberger, *Geophys. J. R. Astron. Soc.* **76**, 399 (1984).
12. This research was supported by NSF grant EAR-8120954 and Japan Society for the Promotion of Science Study Abroad program.

* Present address: Department of Earth Sciences, Nagoya University, Nagoya 464, Japan.

† Present address: Photon Factory, National Laboratory for High-Energy Physics, Oho-Machi, Ibaraki 305, Japan.

‡ Present address: Geophysical Institute, Faculty of Science, University of Tokyo, Tokyo, Japan.

21 November 1983; accepted 14 March 1984

A Candidate Magnetic Sense Organ in the Yellowfin Tuna, *Thunnus albacares*

Abstract. *Single-domain magnetite crystals have been isolated and characterized from tissue located in a sinus within the dermethmoid bone of the skull of the yellowfin tuna, Thunnus albacares. Their chemical composition, narrow size distribution, and distinctive crystal morphology indicate that these crystals are biochemical precipitates. Experiments on the interaction between particles reveal the organization of the particles in situ and suggest a possible form for candidate magnetoreceptor organelles. The consistent localization of such particles with similar arrangement within the dermethmoids of this and other pelagic fishes suggests that the ethmoid region is a possible location for a vertebrate magnetic sense organ.*

Magnetic material has been detected in the tissues of various metazoan species (1-5). Although the material is inferred to be magnetite, in many cases this has not yet been established, and external contaminants have not been excluded as possible sources of magnetic remanence. Even in the homing pigeon and the honey bee, detailed localization of the magnetite has proved difficult to ascertain, and the particles have not been isolated or characterized previously (3, 4). For many of the species studied, behavioral evidence for magnetic sensitivity is lacking or in dispute.

Earlier we reported reproducible conditioned responses to earth-strength magnetic fields in the yellowfin tuna, *Thunnus albacares* (6). We now report the detection, extraction, and characterization of magnetite crystals from tissue within a sinus formed by the dermethmoid bone of the skull of this species. The crystals have a narrow size distribution, are single magnetic domains, and have morphologies similar to other biochemically formed magnetites. Studies of the interactions between particles suggest that the crystals are arranged in groups or chains in the dermethmoid tissue. Magnetite-based magnetoreceptor organelles arranged in vivo in a form consistent with these observations could provide these fish with a sensitive magnetoreception system.

To distinguish magnetic material with a possible magnetoreceptive function from other deposits, we sought to identify a tissue with the following characteristics: (i) it should have a high remanent magnetic moment concentrated in a

small volume of sample compared with other tissues from the same fish; (ii) the anatomical position of the magnetic tissue must be consistent from fish to fish; (iii) the bulk magnetic properties, including particle coercivity, should be similar in different individuals and in different species of fish; and (iv) it should be innervated.

Tissue and organ samples, including bones of the body and the skull, skin, sense organs, viscera, and swimming muscles, were dissected from three 1-year-old yellowfin tuna (fork length, 40 to 50 cm) with glass microtome knives and handled with nonmetallic tools in a magnetically shielded, dust-free clean room. Although subsequent dissections focused on the most magnetic tissue, other samples were measured in all fish. Samples were washed in glass-distilled water, frozen in liquid nitrogen, exposed to strong fields from a cobalt-samarium magnet or an air-core impulse solenoid (7), and tested for isothermal remanent magnetization (IRM) in a superconducting magnetometer. We extracted the magnetic material for other tests by combining the magnetic tissue from several fish, grinding the tissues in a glass tissue grinder, extracting released fats with ether, digesting the remaining cellular material in Millipore filtered 5 percent sodium hypochlorite solution (commercial bleach), and briefly treating the residue with 0.5M EDTA (pH 7.1). After centrifuging and washing, aggregates of black particles could be separated magnetically from the residue; control samples of originally nonmagnetic tissues yielded no such product. The magnetic




Fig. 1. The one-axis Af demagnetization and IRM acquisition curves for the dermethmoids of four yellowfin tuna. Saturation magnetization (1 to 3×10^3 pA·m²) is 100 percent and natural remanent magnetization is 0 percent in both plots. Error bars represent standard errors of the means.

powder extracted from the dermethmoid tissue was analyzed by x-ray diffraction, electron microprobe, and transmission electron microscopy (TEM).

Of 17 tissues and organ samples examined for magnetic remanence, 15 had mean moments less than 500 pA·m², and two (eye tissue and dermethmoid bone) had moments greater than 1000 pA·m² (8). The intensities of magnetization of these samples identified the frontal and dermethmoid bones as the samples containing the greatest concentrations of magnetic material (8). Subdivision and remeasurement of the dermethmoids from a number of fish suggested that the magnetic material was contained in a sinus formed within the dermethmoid bone. Because the dermethmoid bones acquired greater moments (260 to 3000 pA·m²) than the frontal bones (59 to 300 pA·m²) and were always clearly magnetic, we focused our remaining studies on the dermethmoid bone and on the tissue it contained in particular.

The frozen dermethmoid tissues of seven yellowfin tuna had natural remanent magnetization moments at or below the instrument noise level (< 50 pA·m²). We magnetized these samples (600 to 3000 pA·m²) and allowed them to warm to room temperature, measuring their moments at 5-minute intervals. The moments retained by the samples all decayed with time, although not all lost their moments completely within the period of the experiments (1 hour). This observation suggests that, as the tissues thawed, the orientation of the magnetic particles became randomized through thermal agitation.

We washed and refroze the dermethmoids of four fish, subjected them to magnetic fields of progressively increasing strength with the impulse solenoid,

and then demagnetized them with progressively increasing alternating fields (Af). The magnetic moment remaining after each step in these procedures was measured in the magnetometer. The dermethmoids acquired virtually all of their magnetization in fields between 10 and 200 mT and lost it again in alternating fields between 10 and 100 mT (Fig. 1). The absence of the multi-domain magnetite particles detected by Zoeger *et al.* (5) in the Pacific dolphin, *Delphinus delphis*, is indicated by the flatness of the Af demagnetization curve below peak fields of 10 mT. The almost complete saturation of the samples in fields above 200 mT rules out the presence of hematite and metallic iron alloys, which will continue to acquire remanence in fields above 1000 mT.

If the magnetic particles producing the moment were uniformly dispersed throughout the dermethmoid tissue, the IRM acquisition and Af demagnetization curves would be symmetrical about the




Fig. 2. Isolated magnetite crystals from the yellowfin tuna dermethmoid tissue. Scale bar is 100 nm.

50 percent magnetization point. This follows because magnetic moments that are aligned by a given impulse field level should also be moved by an alternating field of the same strength. Interactions between the particles aid Af demagnetization and inhibit IRM acquisition, displacing the curves and causing their intersection to fall below the 50 percent magnetization point (9). However, the abscissa of the intersection point still provides a good estimate of the median coercivity of magnetic particles in the sample (9). The ordinate and the abscissa of the intersection of the Af demagnetization and IRM acquisition curves for the yellowfin tuna dermethmoids were at 30 percent magnetization and 40 mT, respectively (Fig. 1). These data are compatible with the presence of about 8.5×10^7 single-domain magnetite particles in the dermethmoid tissue; these particles are approximately 50 nm in length, have axial ratios of about 0.8 (10), and are organized into interacting groups or chains (9, 11).

An x-ray diffraction pattern identified magnetic particles extracted from the dermethmoid tissue as crystalline magnetite (12). Electron microprobe (Cameca MBX) analysis showed that the crystals were pure, containing no measurable titanium, chromium, or manganese (11). In TEM, the isolated crystals were 45 ± 5 nm in length and 38 ± 5 nm in diameter (mean \pm standard error of the mean) (Fig. 2). These dimensions fall within the single-domain stability field of magnetite (13), and their sizes and axial ratios match the particle coercivities measured in whole tissues. The crystals do not conform to the octahedral crystal morphology or lognormal size-frequency distributions normally shown by geologic or synthetic magnetites (14). Nonoctahedral crystal habits and uniform size distributions are characteristic of chiton and bacterial magnetites (15), which suggests that crystal morphology is a useful means of distinguishing biologic from nonbiologic magnetites (15, 16).

The properties and organization of the magnetite particles in the dermethmoid tissue of the yellowfin tuna meet preconditions for use in magnetoreception and suggest a possible form for magnetite based magnetoreceptors. Their chemical composition, uniform size, and biologically distinctive morphology are evidence of closely controlled biomineralization processes and, consequently, magnetic properties. The crystals will have a coupling energy with the geomagnetic field of about 0.1 kT. They are therefore too small to contribute individually to magnetoreception since their net

alignment, as given by the Langevin function, will be poor (17). Organization of the particles into chains similar to those in the magnetosomes of magnetotactic bacteria (18) will yield greater coupling energies and is consistent with the interactions between the particles detected in the dermethmoid tissue. The decay with warming of the IRM acquired by the dermethmoid tissue indicates that the particle groups are at least partially free to rotate. Taken together, these results suggest an association of the particles with a mechanoreceptor that detects the position or movement of the groups. Theoretical analyses (19) show that chains of 20 to 60 particles would provide ideal coupling energies with the geomagnetic field for use in magnetoreception. Assuming that the 8.5×10^7 particles detected in the dermethmoid tissue are arranged in such a fashion, a magnetite-based magnetoreception system in the yellowfin tuna could resolve magnetic field direction to within a few seconds of arc, or magnetic field intensity differences of 1 to 100 nT (19).

Gross dissection of the dermethmoid region of the yellowfin tuna revealed the supraorbital trunk nerve, which carries branches of the trigeminal, facial, and anterior lateral line nerves and which ramifies in the ethmoid region. Histological studies have suggested the presence of nerve axons in the dermethmoid tissue (20). A suitable physical and possible neural basis for previously demonstrated behavioral responses to magnetic fields has thus been demonstrated for the first time in one species. Our magnetometry results are consistent in phylogenetically distant fishes (12) and, along with similar results for other vertebrates (1, 4, 5), suggest that the ethmoid region of the skull is a likely site for a vertebrate magnetic sense organ.

Note added in proof: Magnetite crystals isolated from the dermethmoid tissue of chinook salmon, *Oncorhynchus tshawytscha*, are organized in chains when viewed in TEM (21).

MICHAEL M. WALKER

Southwest Fisheries Center Honolulu Laboratory, National Marine Fisheries Service, Honolulu, Hawaii 96812, and Department of Zoology, University of Hawaii, Honolulu 96822

JOSEPH L. KIRSCHVINK,

SHIH-BIN R. CHANG

Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena 91125

ANDREW E. DIZON

Southwest Fisheries Center La Jolla Laboratory, National Marine Fisheries Service, La Jolla, California 92038

References and Notes

1. R. R. Baker, J. G. Mather, J. H. Kennaugh, *Nature (London)* **301**, 78 (1983); M. Hanson, L. Karlsson, H. Westerberg, *J. Comp. Biochem. Physiol.*, in press; J. G. Mather and R. R. Baker, *Nature (London)* **291**, 151 (1981); D. Presti and J. D. Pettigrew, *ibid.* **285**, 99 (1980).
2. D. S. Jones and B. J. MacFadden, *J. Exp. Biol.* **96**, 1 (1982).
3. J. L. Gould, J. L. Kirschvink, K. S. Deffeyes, *Science* **201**, 1026 (1978).
4. C. Walcott, J. L. Gould, J. L. Kirschvink, *ibid.* **205**, 1027 (1979).
5. J. Zoeger, J. R. Dunn, M. Fuller, *ibid.* **213**, 892 (1981).
6. M. M. Walker, A. E. Dizon, J. L. Kirschvink, *Oceans 82* (IEEE, Washington, D.C., 1982), pp. 755-758; M. M. Walker, in *Mechanisms of Migration in Fishes*, J. D. McCleave, G. P. Arnold, J. J. Dodson, W. Neill, Eds., in press.
7. J. L. Kirschvink, in *Biomagnetism: An Interdisciplinary Approach*, S. Williamson, G.-L. Romani, L. Kaufman, I. Modena, Eds. (Plenum, New York, 1983), p. 501.
8. Samples with mean moments less than 500 pA·m²: liver, pyloric caecum, intestine, red muscle, white muscle, brain, parietal bone, gill, skin, peduncle tendon, frontal bone, pectoral fin, posterior brain case, dorsal fin, cardiac muscle. Samples with mean moments greater than 1000 pA·m²: eye (1242 ± 526.3 , $N = 4$), dermethmoid bone (1320.6 ± 224.0 , $N = 15$). Background signal in the magnetometer was less than or equal to 50 pA·m². All samples had intensities of magnetization (that is, moments per gram of tissue) less than or equal to 62.5 pT except the frontal bones (44.3, 162.9 pT) and dermethmoid bones (127.0 ± 32.8 pT, $N = 7$).
9. S. Ciszowski, *Phys. Earth Planet. Inter.* **26**, 56 (1981).
10. M. W. McElhinny, *Paleomagnetism and Plate Tectonics* (Cambridge University Press, London, 1973), p. 357.
11. M. M. Walker, A. E. Dizon, J. L. Kirschvink, in *Magnetite Biomineralization and Magnetoreception in Organisms: A New Magnetism*, J. L. Kirschvink, D. S. Jones, B. J. MacFadden, Eds., in preparation.
12. Lattice parameter $a_0 = 0.8358 \pm 0.004$ nm; reference value = 0.8396 nm [*Selected Powder Diffraction Data for Minerals*, compiled by the Joint Committee on Powder Diffraction Standards in cooperation with the American Society for Testing and Materials (and others) (Joint Committee on Powder Diffraction Standards, Swarthmore, Pa., 1974)].
13. R. F. Butler and S. K. Banerjee, *J. Geophys. Res.* **80**, 4049 (1975).
14. R. Løvlie, W. Lowrie, M. Jacobs, *Earth Planet. Sci. Lett.* **15**, 157 (1971); T. Sugimoto and E. Matjević, *J. Colloid. Interface Sci.* **74**, 227 (1980).
15. J. L. Kirschvink and H. A. Lowenstam, *Earth Planet. Sci. Lett.* **44**, 193 (1979); R. B. Frankel, R. P. Blakemore, R. S. Wolfe, *Science* **203**, 1355 (1979).
16. H. A. Lowenstam, *Science* **211**, 1126 (1981); T. Matsuda, J. Endo, N. Osakabe, A. Tonomura, T. Arai, *Nature (London)* **302**, 411 (1983); K. M. Towle and T. T. Moench, *Earth Planet. Sci. Lett.* **52**, 213 (1981).
17. R. B. Frankel and R. P. Blakemore, *J. Magnet. Magn. Mater.* **15-18**, 1562 (1980); J. L. Kirschvink, *BioSystems* **14**, 193 (1981).
18. D. L. Balkwill, D. Maratea, R. P. Blakemore, *J. Bacteriol.* **141**, 1399 (1980).
19. J. L. Kirschvink and J. L. Gould, *BioSystems* **13**, 181 (1981); J. L. Kirschvink and M. M. Walker, in *Magnetite Biomineralization and Magnetoreception in Organisms: A New Magnetism*, J. L. Kirschvink, D. S. Jones, B. J. MacFadden, Eds. (Plenum, New York, in press); E. D. Yorke, *J. Theor. Biol.* **77**, 101 (1979); *ibid.* **89**, 533 (1981).
20. M. M. Walker, unpublished data.
21. J. L. Kirschvink *et al.*, in preparation.
22. We thank H. A. Lowenstam, K. A. Peterson (Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena 91125), and A. Perry for helpful discussions, critical reviews, and assistance with experiments. C. E. Helsley, B. H. Keating, L. C. Ming, and M. O. Garcia (Hawaii Institute of Geophysics, University of Hawaii, Honolulu 96822) provided use of paleomagnetic laboratory facilities and assistance with x-ray diffraction and electron microprobe analyses. J. P. Revel (California Institute of Technology) made facilities available for electron microscopy. This research was supported in part by a graduate study award from the East-West Center, Honolulu, Hawaii 96848 (M.M.W.), NSF grants PCM82-03627 and BNS83-00301 (J.L.K.), and NOAA funding. Contribution 4003 from the Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, Calif. 91125.

8 August 1983; accepted 26 January 1984

Progressive Accumulation of Toxic Metabolite in a Genetic Leukodystrophy

Abstract. *Progressive accumulation of a cytotoxic metabolite, galactosylsphingosine (psychosine), was found in the brain of the twitcher mouse, a mutant caused by genetic deficiency of galactosylceramidase. Similar abnormal accumulation was also found in the brain of the genetic galactosylceramidase deficiency disease in the dog and in human patients (globoid cell leukodystrophy or Krabbe disease). Galactosylsphingosine was absent in the brains of normal and heterozygous mice. The finding provides support for the psychosine hypothesis as the biochemical pathogenetic mechanism of globoid cell leukodystrophy. Analogous mechanisms may be important in the pathogenesis of other genetic lysosomal diseases.*

Genetic galactosylceramidase deficiency (globoid cell leukodystrophy or Krabbe disease) is known in several mammalian species, including the human (1), the dog (2), and the mouse (3). It is a rapidly progressive fatal disorder with clinical and pathological manifestations almost exclusively restricted to the nervous system, particularly to the white matter and the peripheral nerves. Conceptually the disease is related to the lysosomal storage disorder of Hers (4). Paradoxically, abnormal accumulation of galactosylceramide, the natural sub-

strate of the missing enzyme, does not occur in the nervous system despite the genetic catabolic block (2, 5, 6).

Miyatake and Suzuki (7) in 1972 proposed a hypothesis for the biochemical mechanism that leads to the absence of galactosylceramide accumulation and to the devastating pathology of the white matter, particularly the rapid and total degeneration of the oligodendroglia. The hypothesis has become known as psychosine hypothesis, because a toxic metabolite, galactosylsphingosine (psychosine), which is also a substrate of galac-