



Draft Genome Sequence of *Chloracidobacterium* sp. CP2_5A, a Phototrophic Member of the Phylum *Acidobacteria* Recovered from a Japanese Hot Spring

Lewis M. Ward,^a Shawn E. McGlynn,^b Woodward W. Fischer^a

Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, California, USA^a;
Earth-Life Science Institute, Tokyo Institute of Technology, Meguro-ko, Japan^b

ABSTRACT The phylum *Acidobacteria* contains a single known phototrophic member, *Chloracidobacterium thermophilum*, which was recovered from a hot spring metagenome from Yellowstone National Park. Here, we expand the diversity of the genus *Chloracidobacterium* with a genome recovered from a hot spring in Japan, extending the known range of this lineage to a new continent.

The phylum *Acidobacteria* comprises widespread but enigmatic bacteria; many subdivisions have been observed in environmental sequencing data, but only a few cultured or well-characterized representatives exist (1). The only known phototroph in this phylum, *Chloracidobacterium thermophilum*, was originally recovered from metagenomic data from a Yellowstone National Park hot spring (2) and was recently isolated (3). Here, we report the draft genome of *Chloracidobacterium* sp. strain CP2_5A, recovered from a metagenome of a Japanese hot spring.

The metagenome from which this genome was recovered was described in detail by Ward (4). In brief, CP2_5A was recovered from a cone-forming microbial mat from a moderately sulfidic, alkaline hot spring at Nakabusa Onsen in Nagano prefecture, Japan. At the time of sampling, the temperature at the collection site was 32°C with a pH measurement of 8.3.

Cells were lysed and DNA was preserved in the field using Zymo Terralyzer BashingBead Matrix and Xpedition Lysis Buffer. DNA was extracted and purified with a Zymo Soil/Fecal DNA extraction kit (Zymo Research, Irvine, CA, USA) and quantified with a Qubit version 3.0 fluorimeter (Life Technologies, Inc., Carlsbad, CA, USA). Purified DNA was submitted to SeqMatic LLC (Fremont, CA, USA) for library preparation and sequencing via Illumina HiSeq technology. Raw sequences were assembled with MegaHit version 1.02 (5) and binned using MetaWatt version 3.5.2 (6). The genome was uploaded to RAST (7) for overall characterization and was assessed for completeness and contamination using CheckM (8).

The CP2_5A genome bin is estimated to be 92% complete as determined by CheckM. The genome is 3.41 Mb, with a GC content of 64.2%, and comprises 385 contigs containing 3,023 coding sequences and 52 RNAs. The 16S gene recovered from CP2_5A is 98% identical to that of *C. thermophilum* strain E. The CP2_5A genome contains a complete suite of genes for anoxygenic phototrophy, including a type 1 reaction center (RCI) and a largely complete bacteriochlorophyll synthesis pathway (only *bchM* was not recovered in the genome bin). Consistent with the photoheterotrophic lifestyle of the isolated *C. thermophilum* strain, CP2_5A does not encode carbon fixation pathways. The CP2_5A genome encodes two *bc* complexes, an alternative complex III, a low-O₂-affinity A-family heme-copper oxidoreductase, and a *bd*

Received 3 July 2017 Accepted 18 August 2017 Published 5 October 2017

Citation Ward LM, McGlynn SE, Fischer WW. 2017. Draft genome sequence of *Chloracidobacterium* sp. CP2_5A, a phototrophic member of the phylum *Acidobacteria* recovered from a Japanese hot spring. Genome Announc 5:e00821-17. <https://doi.org/10.1128/genomeA.00821-17>.

Copyright © 2017 Ward et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Lewis M. Ward, lward@caltech.edu.

cytochrome *c*. The RCI of CP2_5A is similar to that of the *Chloracidobacterium* isolate: it branches basal to those of phototrophic *Chlorobi* spp. but more derived than the simple RCIs found in *Heliobacteria* spp. While one of the *bc* complexes in CP2_5A clusters with that of *C. thermophilum* strain E, the other is instead closely related to that from the recently discovered phototrophic phylum *Gemmatimonadetes* (9), potentially marking horizontal gene transfer between these two clades.

The degree of genomic and metabolic similarity between *Chloracidobacterium* strains found in hot springs across at least two continents suggests a broadly conserved ecology between microbial lineages in similar geochemical environments across large spatial scales.

Accession number(s). This whole-genome shotgun project was deposited in DDBJ/EMBL/GenBank under the accession number [NKPT00000000](https://doi.org/10.1128/genomeA.00897-14).

ACKNOWLEDGMENTS

L.M.W. acknowledges support from NASA NESSF (grant NNX16AP39H), NSF (grant OISE 1639454), NSF GROW (grant DGE 1144469), the Lewis and Clark Fund for Exploration and Field Research in Astrobiology, and the ELSI Origins Network. S.E.M. acknowledges support from a MEXT KAKENHI grant-in-aid for challenging exploratory research (grant award 15K14608). W.W.F. acknowledges the support of NASA Exobiology award NNX16AJ57G, the David and Lucile Packard Foundation, and a Stanford University Blaustein Fellowship.

We thank Katsumi Matsura and the Environmental Microbiology Laboratory at Tokyo Metropolitan University for laboratory support.

REFERENCES

1. Stamps BW, Corsetti FA, Spear JR, Stevenson BS. 2014. Draft genome of a novel *Chlorobi* member assembled by tetranucleotide binning of a hot spring metagenome. *Genome Announc* 2(5):e00897-14. <https://doi.org/10.1128/genomeA.00897-14>.
2. Bryant DA, Costas AMG, Maresca JA, Chew AGM, Klatt CG, Bateson MM, Tallon LJ, Hostetler J, Nelson WC, Heidelberg JF, Ward DM. 2007. *Candidatus Chloracidobacterium thermophilum*: an aerobic phototrophic acidobacterium. *Science* 317:523–526. <https://doi.org/10.1126/science.1143236>.
3. Tank M, Bryant DA. 2015. *Chloracidobacterium thermophilum* gen. nov., sp. nov.: an anoxygenic microaerophilic chlorophotoheterotrophic acidobacterium. *Int J Syst Evol Microbiol* 65:1426–1430. <https://doi.org/10.1099/ijs.0.000113>.
4. Ward LM. 2017. Microbial evolution and the rise of oxygen: the roles of contingency and context in shaping the biosphere through time. Ph.D. dissertation, California Institute of Technology, Pasadena, CA. <https://doi.org/10.7907/Z9BZ642S>.
5. Li D, Luo R, Liu CM, Leung CM, Ting HF, Sadakane K, Yamashita H, Lam TW. 2016. MEGAHIT v1.0: a fast and scalable metagenome assembler driven by advanced methodologies and community practices. *Methods* 102: 3–11. <https://doi.org/10.1016/j.ymeth.2016.02.020>.
6. Strous M, Kraft B, Bisdorf R, Tegetmeyer HE. 2012. The binning of metagenomic contigs for microbial physiology of mixed cultures. *Front Microbiol* 3:410. <https://doi.org/10.3389/fmicb.2012.00410>.
7. Aziz RK, Bartels D, Best AA, Dejongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, Mcneil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
8. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
9. Zeng Y, Feng F, Medová H, Dean J, Koblížek M. 2014. Functional type 2 photosynthetic reaction centers found in the rare bacterial phylum *Gemmatimonadetes*. *Proc Natl Acad Sci U S A* 111:7795–7800. <https://doi.org/10.1073/pnas.1400295111>.